

(2'-deoxy)adenosine + ATP => (d)AMP + ADP (ADK)

D'Eustachio, P.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 77


This document contains 1 reaction ([see Table of Contents](#))

(2'-deoxy)adenosine + ATP => (d)AMP + ADP (ADK) ↗

Stable identifier: R-HSA-109624

Type: transition

Compartments: cytosol



Cytosolic adenosine kinase (ADK) catalyzes the reactions of adenosine and deoxyadenosine with ATP to yield the corresponding nucleotide monophosphates and ADP (Andres and Fox 1979). The enzyme is substantially more active on adenosine than deoxyadenosine in vitro (Hurley et al. 1985) though studies of cultured cells suggest that both reactions may be physiologically relevant (Hershfield et al. 1982). The enzyme is a monomer complexed with magnesium (Mathews et al. 1998).

Literature references

Hurley, MC., Lin, B., Fox, IH. (1985). Regulation of deoxyadenosine and nucleoside analog phosphorylation by human placental adenosine kinase. *J Biol Chem*, 260, 15675-15681. ↗

Andres, CM., Fox, IH. (1979). Purification and properties of human placental adenosine kinase. *J Biol Chem*, 254, 11388-11393. ↗

Hershfield, MS., Fetter, JE., Small, WC., Bagnara, AS., Williams, SR., Ullman, B. et al. (1982). Effects of mutational loss of adenosine kinase and deoxycytidine kinase on deoxyATP accumulation and deoxyadenosine toxicity in cultured CEM human T-lymphoblastoid cells. *J Biol Chem*, 257, 6380-6386. ↗

Mathews, II., Erion, MD., Ealick, SE. (1998). Structure of human adenosine kinase at 1.5 Å resolution. *Biochemistry*, 37, 15607-20. ↗

Editions

2010-02-06

Revised

D'Eustachio, P.